



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number : 0 552 060 A1

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number : 93300278.4

(22) Date of filing : 15.01.93

(51) Int. Cl.<sup>5</sup> : C07D 233/84, C07D 233/90,  
C07D 233/60, C07D 401/06,  
C07D 403/06, C07D 413/06,  
C07D 417/06, A61K 31/35,  
A61K 31/415, A61K 31/395,  
A61K 31/42, A61K 31/425,  
A61K 31/435, A61K 31/44,  
A61K 31/495, A61K 31/50,  
A61K 31/505, C07D 233/54

(30) Priority : 16.01.92 JP 5577/92

(43) Date of publication of application :  
21.07.93 Bulletin 93/29

(84) Designated Contracting States :  
AT BE CH DE DK ES FR GB GR IE IT LI LU MC  
NL PT SE

(71) Applicant : SHIONOGI & CO., LTD.  
1-8, Doshomachi 3-chome, Chuo-ku  
Osaka-shi, Osaka-fu (JP)

(72) Inventor : Sugimoto, Hirohiko  
5-7-7 Senriyama-nishi  
Suita-shi, Osaka-fu (JP)


Inventor : Ogata, Masaru  
8-20-8 Sumiyoshi-yamate, Higashinada-ku  
Kobe-shi, Hyogo-ken (JP)  
Inventor : Matsumoto, Hiroshi  
5-10-25 Yamatedai

Ibaraki-shi, Osaka-fu (JP)  
Inventor : Sugita, Ken-ichi  
2-3-53 Kokubu-nishi  
Kashiwara-shi, Osaka-fu (JP)  
Inventor : Sato, Akihiko  
2-3-21 Minami-mimihara  
Ibaraki-shi, Osaka-fu (JP)  
Inventor : Fujiwara, Tamio  
2-21-107 Oogi, Higashinada-ku  
Kobe-shi, Hyogo-ken (JP)

(74) Representative : Hardisty, David Robert et al  
BOULT, WADE & TENNANT 27 Farnival Street  
London EC4A 3DF (GB)

(54) Imidazole derivatives having anti-HIV activity.

(57) A novel imidazole derivatives of formula I :



wherein R<sup>1</sup> is hydrogen, alkyl, halogen, or optionally substituted aryl; R<sup>2</sup> is alkyl, optionally substituted aryl optionally substituted aralkyl, or optionally substituted hetero ring group; R<sup>3</sup> is hydrogen, alkyl, optionally substituted aryl, optionally substituted aralkyl, or optionally substituted hydroxyalkyl; R<sup>4</sup> is hydrogen, alkyl, halogen, acyl, optionally substituted hydroxyalkyl, optionally esterified or amidated carboxyl group, hydroxy group, aryl or arylthio; X is S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or Se; n is an integer of 1 to 3, or a pharmaceutically acceptable salt thereof, said derivative having anti-HIV activity and being useful for the treatment of HIV infections.

EP 0 552 060 A1

The present invention relates to novel imidazole derivatives, more particularly it relates to novel imidazole derivatives having anti-retrovirus activity and pharmaceutical compositions effective for the treatment of retrovirus infections such as human immunodeficiency syndromes.


AIDS (acquired immunodeficiency syndrome) is caused by a member of retrovirus, human immunodeficiency virus (HIV). In spite of great efforts made by researchers there have been no effective methods for treating AIDS so far, and the disease is becoming a more and more serious problem.

It has been reported that pyrimidine nucleoside compounds can be useful anti-retrovirus agents because of the inhibitory activity against the reverse transcription of RNA of retroviruses and many compounds have been synthesized. Among them, azidodeoxythymidine (AZT) has been practically used for treating AIDS. AZT, however, cannot be used broadly due to its high toxicity. Accordingly, the development of anti-HIV compounds which are highly effective and less toxic has been demanded strongly.

It has been reported that acyclovir, an acyclonucleoside derivative, has an inhibitory effect on herpes virus and hence a lot of acyclonucleosides with various side chains were synthesized [J. Heterocyclic Chem. 23, 289 (1986)]. The literature failed to mention about activities of such compounds against retrovirus.

The present inventors have made an intensive investigation for the purpose of developing compounds having anti-retrovirus activity and being useful for the treatment of AIDS; they found that certain imidazole derivatives have such an activity and can prevent HIV infection.

Thus, the present invention provides compounds of the formula I:



wherein R<sup>1</sup> is hydrogen, alkyl, halogen, or optionally substituted aryl; R<sup>2</sup> is alkyl, optionally substituted aryl, optionally substituted aralkyl, or optionally substituted hetero ring group; R<sup>3</sup> is hydrogen, alkyl, optionally substituted aryl, optionally substituted aralkyl, or optionally substituted hydroxyalkyl; R<sup>4</sup> is hydrogen, alkyl, halogen, acyl, optionally substituted hydroxyalkyl, optionally esterified or amidated carboxyl group, hydroxy group, aryl or arylthio; X is S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or Se; n is an integer of 1 to 3, or a pharmaceutically acceptable salt thereof.

For purposes of the present invention, as disclosed and claimed herein, the following terms are defined below.

The term "alkyl" means a straight or branched saturated hydrocarbon radical having one to ten carbon atoms, preferably, a straight or branched alkyl radical having one to six carbon atoms. Examples of alkyl include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, neopentyl, s-pentyl, t-pentyl, n-hexyl, neohexyl, i-hexyl, s-hexyl, t-hexyl and the like.

For purposes of the present invention, a straight or branched lower alkyl having one to four carbon atoms such as methyl, ethyl, n-propyl, isopropyl or t-butyl is particularly preferred.

The term "halogen" means fluorine, chlorine, bromine and iodine.

The term "acyl" means an aromatic or aliphatic acyl group. Examples of aromatic acyl groups are benzoyl, 4-nitrobenzoyl, 4-tert-butylbenzoyl, benzenesulfonyl, toluenesulfonyl and the like. Examples of aliphatic acyl groups are formyl, acetyl, propionyl, butyryl, valeryl and the like. Aliphatic acyl group is preferred, and formyl is the most preferable group.

The term "aryl" means an aromatic ring group having six to twelve carbon atoms. Examples of aryl group include phenyl, tolyl, xylyl, biphenyl, naphthyl and the like, with the preference of phenyl.

In the definition of "optionally substituted aryl", examples of the substituents include alkyl, halogen, nitro, amino, hydroxy, alkoxy, and aralkyloxy as defined above, with the preference of methyl, fluorine, and chlorine.

The term "hydroxyalkyl" means those derived from an alkyl group as defined above. Preferred hydroxyalkyl groups are hydroxymethyl and hydroxyethyl.

In the definition of "optionally substituted hydroxyalkyl", substituted hydroxyalkyl means those which are derived by substituting an alkyl, aryl, aralkyl or acyl for the hydrogen of the hydroxy group, namely, it includes alkoxyalkyl, aryloxyalkyl, aralkyloxyalkyl, acyloxyalkyl and the like. Especially, benzyloxyalkyls are preferable.

Examples of "optionally esterified carboxyl group" include carboxyl group, methoxycarbonyl group, ethoxycarbonyl, tert-butoxycarbonyl, benzyloxycarbonyl and the like, with the preference of ethoxycarbonyl.

Salts of compounds of the formula (I) are pharmaceutically acceptable salts formed between a compound


(I) and an inorganic acid such as hydrochloric acid, nitric acid, phosphoric acid, sulfuric acid, hydrobromic acid, hydroiodic acid, nitrous acid, phosphorous acid or the like; or a non-toxic organic acid such as aliphatic mono- or dicarboxylic acid, phenyl-substituted alkanolic acid, hydroxy-mono- or hydroxy-di-alkanoic acid, aromatic acid, aliphatic or aromatic sulfonic acid or the like. Examples of pharmaceutically acceptable acid addition salts include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, nitrate, phosphate, secondary phosphate, primary phosphate, metaphosphate, pyrophosphate, hydrochloride, hydrobromide, hydroiodide, hydrofluoride, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caproate, heptanoate, propionate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, mandelate, butyn-1,4-dioate, hexyn-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, phthalate, telephthalate, benzenesulfonate, toluenesulfonate, chlorobenzenesulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate,  $\beta$ -hydroxybutyrate, glycolate, malate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate and the like. For purposes of the invention, inorganic salts are preferred, and hydrochloride and nitrate are particularly preferable.

All the compounds (I) as defined above commonly possess the objective anti-retrovirus activity and are useful for the treatment of AIDS. There are certain preferable compounds which have higher anti-HIV activity, e.g., compounds (I) wherein  $R^1$  is hydrogen,  $C_1$ - $C_3$  alkyl, in particular, methyl, ethyl or propyl, halogen, in particular, bromine, or phenyl;  $R^2$  is  $C_1$ - $C_4$  alkyl, in particular, butyl, phenyl, phenyl substituted by halogen or lower alkyl, in particular, dimethylphenyl;  $R^3$  is  $C_1$ - $C_3$  alkyl, in particular, ethyl, benzyl, or hydroxyalkyl, in particular, hydroxyethyl;  $R^4$  is hydrogen,  $C_1$ - $C_3$  alkyl, in particular, methyl or ethyl, halogen, in particular, bromine, formyl, hydroxymethyl, ethoxycarbonyl, carbamoyl, benzyloxymethyl, hydroxyl or phenylthio group; and X is S or  $CH_2$  are preferred for purposes of the invention.

As can be understood easily from the formula I, the compounds (I) of the present invention can be in two position isomers in which a substituent of the formula  $-CH_2-O-R^3$  is attached to either of nitrogen atoms in the imidazole ring.

The preparation of the compounds (I) of the present invention can be carried out by any of known methods, for example, those shown by the synthetic routes A, B, C, D, E, F, G, H, and/or I below.

#### Synthetic Route A



In the reaction scheme above,  $X'$  means S or Se and Y means halogen.

The reactions in the synthetic route A can be carried out according to the teachings shown at every step by using an appropriate imidazole derivative, e.g. 4-substituted-imidazole as the starting material. Each reaction can be carried out by employing conditions known to those skilled in the art.

For example, in the first step, an iodine atom is introduced at the 5-position of imidazole ring in the presence of a base in a solvent such as methylene chloride or the like. In the second step, the resultant imidazolyl iodide is added to a mixture of sodium hydride and a solution of thiophenol (or selenophenol) in an appropriate solvent such as dry dimethylformamide to introduce a phenylthio (or phenylseleno) group at the 5-position of imidazole ring. In the third step, a desired substituent,  $-(CH_2)_n-O-R^3$ , is introduced at a nitrogen atom on imidazole ring. This reaction tends to give a mixture of position isomers as a product, in which the group  $-(CH_2)_n-O-R^3$  is attached to either nitrogen atoms at 1- or 3-positions on imidazole ring. The isomer having the substituent at the desired position can be isolated by any of conventional methods such as silica gel chromatography. In the fourth step, the imidazole derivatives obtained in steps 1 to 3 above are converted into the objective compound (I). Thus, the hydrogen at the 2-position of imidazole ring is replaced by a desired substituent in the presence


of an electrophilic reagent known per se in the art, for instance, n-butyllithium.

### Synthetic Route B

5

10

15




20 In the reaction scheme above, X' and Y are the same as defined in the Route A. The process is carried out substantially in the same manner as in Route A, except that the substituent R<sup>4</sup> is introduced before the introduction of -CH<sub>2</sub>-O-R<sup>3</sup> to the imidazole ring.

### Synthetic Route C

25

30

35




40 In the reactions scheme above, X' and Y are the same as defined in Routes A and B. This process is carried out substantially in the same manner as in Route A or B, using 2,4-substituted-imidazoles as starting materials. Most of the starting materials are known compounds or, if necessary, they may be readily prepared by conventional reactions.

### Synthetic Route D

45

50

55



X=S, SO, SO<sub>2</sub>, Se or CH<sub>2</sub>


In the reactions scheme above, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, and X are the same as defined before.

The starting material A, an α-acylamino-α-substituted-ketone, is refluxed in an organic acid solvent such

as acetic acid in the presence of ammonia or the salts with organic acids or inorganic acids to give an objective 2,4,5-substituted imidazole. The subsequent reaction may be carried out according to the manner in Route A, B, or C.


The starting material A can be prepared, for example, through either of the following routes.

1)  $X = CH_2$




Amino acid is halogenated with thionyl halide or phosphorus oxychloride and then refluxed with alcohol to give the ester, which is reacted with an acid anhydride or an activated organic acid such as acid halide in the presence of a base to give an N-acetylamino acid ester. This is treated with a carbanion reagent as Grignard reagents or organic lithium reagent in an aprotic solvent to give the starting material A.

2)  $X = CH_2$




Acetylaminoacetic acid diester is in the presence of a base alkylated with a substituted methyl halide in an aprotic solvent to give  $\alpha$ -aminoacyl- $\alpha$ -substituted malonic acid diester, which is hydrolyzed and decarboxylated, followed by esterification. The resulting ester is treated in the same manner as aforementioned to give the starting material A.

3)  $X = S, SO, SO_2$  or Se







A substituted ( $\text{XR}^2$ ) acetic acid ester is treated with azide transfer reagent in the presence of a base in an aprotic solvent at a temperature of from  $-100^\circ\text{C}$  to room temperature. The azide transfer agent includes, for example, benzenesulfonylazide, 2,4,6-trimethylbenzenesulfonylazide, diphenylphosphorylazide and the like. The reaction mixture is subjected to the same reaction as shown above to give the corresponding ketone. The azide group of the ketone is reduced and acylated to give the starting material A.

Synthetic Route E

15 Compounds (I) wherein X is SO or SO<sub>2</sub> can be readily obtained by oxidation of the corresponding sulfides. The oxidation is carried out in an appropriate solvent such as methylene chloride, acetonitrile and the like solvent with one or two mole of an appropriate oxidizing agent such as sodium periodate, m-chloroperbenzoic acid, hydrogen peroxide, sodium dichromate, potassium permanganate, chromic acid and the like.

20 Thus obtained compounds (I) in the synthetic route A, B, C, D or E above may be further modified according to the synthetic route F, G, H, or I as described below to yield desired Imidazole derivatives.

Synthetic Route FSynthetic Route GSynthetic Route H



Although the present invention is by no means restricted to the production of a compound (I), there are a certain preferable synthetic route for each compound depending on the definition of "X" in the formula (I) as shown below.

Preferable synthetic routes of compounds (I)	
Synthetic route	X of formula (I)
A, B, C	S, Se
D	S, SO, SO <sub>2</sub> , Se, CH <sub>2</sub>
F	S, Se, CH <sub>2</sub>
G, H, I	S, SO, SO <sub>2</sub> , Se, CH <sub>2</sub>

The compounds of the present invention can be orally or parenterally administered. For oral administration, the compounds of the present invention can be used in any formulations in conventional forms such as solid formulations including tablets, powders, granules, capsules and the like; aqueous solutions; oily suspensions; and liquid formulations including syrups, elixirs and the like. For parenteral administration, the compounds of the present invention can be formulated as an aqueous or oily suspension injections. The formulations of the invention can be prepared by using conventional adjuvant such as excipients, binders, lubricants, aqueous solvents, oily solvents, emulsifiers, suspending agents or the like. The formulations may comprise further supplements such as preservatives, stabilizers or the like.

The following example is set forth to further describe the invention but is in no way meant to be construed as limiting the scope thereof.


Example 11-Benzoyloxymethyl-4-ethyl-2-methyl-5-phenylthioimidazole

5

10

15

20

(1) 2-Methyl-4-ethyl-5-iodoimidazole 1

25

30

To a solution of 910 mg of NaOH (23 mmol) in 44 ml of water are added 38 ml of methylene chloride and 2.5 g of 2-methyl-4-ethylimidazole (23 mmol). A solution of 5.8 g of iodine (23 mmol) in 44 ml of methylene chloride and 22 ml of methanol is dropwise added under ice cooling. Fifteen minutes later, the mixture is extracted with methylene chloride. The extract is washed with water, dried over  $Na_2SO_4$ , filtered, and the filtrate is concentrated. The resulting crude product is chromatographed on a column of silica gel, eluting with 3% methanol/methylene chloride to give 3.29 g of the desired 2-methyl-4-ethyl-5-iodoimidazole 1. Yield, 62%; mp. 156 - 159°C

1:
 $^1H-NMR(CDCl_3-TMS)\delta ppm$ : 1.19(t, J=7.5Hz, 3H), 2.39(s, 3H), 2.57(q, J=7.5Hz, 2H), 9.05(br, 1H)

35

(2) 2-Methyl-4-ethyl-5-phenylthioimidazole 2

40

To a solution of 2.24 g of thiophenol (20.3 mmol) in 16 ml of dry dimethylformamide is added 1.4 g of sodium hydride (60% oil suspension: 35 mmol) under ice cooling. Ten minutes later, 3.2 g of 2-methyl-4-ethyl-5-iodoimidazole (13.6 mmol) is added to the mixture, which is heated at 130°C in a nitrogen atmosphere. Two hours later, the dimethylformamide is evaporated in vacuo. The residue is mixed with dry ice and extracted with methylene chloride. The extract is washed with water, dried over  $Na_2SO_4$ , filtered and the filtrate is concentrated in vacuo. The crude product is chromatographed on a column of silica gel, eluting with 3% methanol/methylene chloride. The product is recrystallized from ethyl acetate/isopropyl ether to give 1.9 g of 2-methyl-4-ethyl-5-phenylthioimidazole 2. Yield, 64%; mp. 171°C

45

2:
 $^1H-NMR(CDCl_3-TMS)\delta ppm$ : 1.19(t, J=7.5Hz, 3H), 2.39(s, 3H), 2.65(q, J=7.5Hz, 2H), 5.40(br, 1H), 7.03-7.26(m, 5H)

50

(3) 1-Benzoyloxymethyl-5-ethyl-2-methyl-4-phenylthioimidazole 3 and 1-benzoyloxymethyl-4-ethyl-2-methyl-5-phenylthioimidazole (1-1)

55

To a solution of 800 mg of 2-methyl-4-ethyl-5-phenylthioimidazole (3.7 mmol) in 8 ml of dry dimethylformamide is added 220 mg of 60% oil suspension of sodium hydride (5.5 mmol) under ice cooling. Ten minutes later, 860 mg of benzoyloxymethyl chloride (5.6 mmol) is added dropwise. Fifteen minutes later, ice water is added to the mixture, which is extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated in vacuo. The residue containing both the non-polar and polar products is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1).

From the former elution is obtained 160 mg of 1-benzoyloxymethyl-5-ethyl-2-methyl-4-phenylthioimidazole 3 as an oil. Yield, 13%.



3:  
<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.09(t,J=7.4Hz,3H), 2.43(s,3H), 2.73(q,J=7.4Hz), 4.52(s,2H), 5.26(s,2H), 7.00-7.31(m,10H)


From the later elution is obtained 810 mg of 1-benzyloxymethyl-4-ethyl-2-methyl-5-phenylthioimidazole (I-1) as an oil. Yield, 65%. The product, when recrystallized from n-hexane, gives crystals melting at 50 - 51°C.

I-1:  
<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.24(t,J=7.4Hz,3H), 2.53(s,3H), 2.69(q,J=7.4Hz,2H), 4.36(s,2H), 5.30(s,2H), 6.97-7.31(m,10H)

Elemental analysis (C <sub>20</sub> H <sub>22</sub> N <sub>2</sub> OS)				
Calcd. (%):	C, 70.97:	H, 6.55:	N, 8.28:	S, 9.47
Found (%):	C, 70.91:	H, 6.58:	N, 8.54:	S, 9.38

### Example 2

#### 1-Ethoxymethyl-4-ethyl-2-methyl-5-phenylthioimidazole (I-2)



To a solution of 500 mg of 4-ethyl-2-methyl-5-phenylthioimidazole 2 (2.3 mmol) obtained in Example 1 (2) in 5 ml of dry dimethylformamide is added 140 mg of 60% oil suspension of sodium hydride (3.5 mmol) under ice cooling. Five minutes later, 330 mg of ethoxycarbonyl chloride (3.5 mmol) is added to the mixture, which is stirred for 15 minutes. The reaction mixture is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (2:1) to give 410 mg of the desired product I-2 as an oil. Yield, 65%.


I-2:  
<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.04(t,J=7.0Hz,3H), 1.23(t,J=7.4Hz,3H), 2.52(s,3H), 2.68(q,J=7.4Hz,2H), 3.35 (q,J=7.0Hz,2H), 5.22(s,2H), 6.98-7.27(m,5H)

Elemental analysis (C <sub>16</sub> H <sub>20</sub> N <sub>2</sub> OS 1/6H <sub>2</sub> O)				
Calcd. (%):	C, 64.49:	H, 7.22:	N, 10.03:	S, 11.48
Found (%):	C, 64.55:	H, 7.37:	N, 10.03:	S, 11.26

### Examples 3 - 18

In Examples 3 to 18, compounds (I-3) - (I-18) are obtained by conducting the reaction substantially in the same manner as that described in Examples 1 and 2. The resultant compounds are summarized in Table 1 below.

Table 1



Exam. No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)salt	Comp. No.
3	S	-CH <sub>3</sub>		-CH <sub>2</sub> -	H	127-128 HCl	I- 3
4	S	-C <sub>2</sub> H <sub>5</sub>		-CH <sub>2</sub> -	H	129-130 HNO <sub>3</sub>	I- 4
5	S	H		-CH <sub>2</sub> -	H	145-146 HNO <sub>3</sub>	I- 5
6	S			-CH <sub>2</sub> -	H	oil	I- 6
7	S	-CH <sub>3</sub>		-C <sub>2</sub> H <sub>5</sub>	H	79-80 HNO <sub>3</sub>	I- 7
8	S	-CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>3</sub> CH <sub>3</sub>	-CH <sub>2</sub> -	H	124-126 HNO <sub>3</sub>	I- 8
9	S	-(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>		-CH <sub>2</sub> -	H	121-122 HNO <sub>3</sub>	I- 9
10	S			-CH <sub>2</sub> -	Br	oil	I-10
11	S	-CH <sub>3</sub>		-C <sub>2</sub> H <sub>5</sub>	Br	90 HNO <sub>3</sub>	I-11
12	S	-C <sub>2</sub> H <sub>5</sub>		-CH <sub>2</sub> -	Br	53-54	I-12
13	S	-(CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>		-CH <sub>2</sub> -	Br	131-132 HCl	I-13
14	S	-C <sub>2</sub> H <sub>5</sub>		-CH <sub>2</sub> -	-CH <sub>3</sub>	oil	I-14
15	S	-C <sub>2</sub> H <sub>5</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	I-15
16	S			-CH <sub>2</sub> -	-CH <sub>3</sub>	oil	I-16
17	S			-CH <sub>2</sub> -	-CH <sub>3</sub>	92.5~93.5°	I-17
18	S			-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	107~108° HCl	I-18

Physicochemical properties of each compound as described in Table 1 are provided below.

I-3:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: (free base) 2.36(s,3H), 4.37(s,2H), 5.32(s,2H), 6.97-7.31(m,10H), 7.82(s,1H)

Elemental analysis ( $\text{C}_{16}\text{H}_{16}\text{N}_2\text{OS HCl}$ )					
Calcd.(%):	C,62.33:	H,5.52:	Cl,10.22:	N,8.06:	S,9.24
Found (%):	C,62.07:	H,5.59:	Cl,10.24:	N,8.02:	S,9.03

I-4:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: 1.25(t,J=7.5Hz,3H), 2.72 (q,J=7.5Hz,2H), 4.38(s,2H), 5.32(s,2H), 6.97-7.31(m,10H), 7.82(s,1H)

Elemental analysis ( $\text{C}_{16}\text{H}_{20}\text{N}_2\text{OS HNO}_3$ )				
Calcd.(%):	C,58.90:	H,5.46:	N,10.85:	S,8.28
Found (%):	C,58.88:	H,5.55:	N,10.93:	S,8.35

I-5:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: (free base) 4.36(s,2H), 5.35(s,2H), 7.07-7.32(m,10H), 7.47(s,1H), 7.89(s,1H)

I-6:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: 4.50(s,2H), 5.27(s,2H), 7.10-7.44 (m,15H), 7.82(s,1H)

I-7:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: (free base) 1.02(t,J=7.0Hz,3H), 2.34(s,3H), 3.35(q,J=7.0Hz,2H), 5.26(s,2H), 6.95-7.27 (m,5H), 7.79(s,1H)

Elemental analysis ( $\text{C}_{13}\text{H}_{16}\text{N}_2\text{OS HNO}_3$ )				
Calcd.(%):	C,50.15:	H,5.50:	N,13.50:	S,10.30
Found (%):	C,50.09:	H,5.50:	N,13.61:	S,10.22

I-8:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: 0.86(t,J=7.5H,3H), 1.41(m,4H), 2.33(s,3H), 2.59(t,J=7.5Hz,2H), 4.48(s,2H), 5.40(s,2H), 7.26-7.35(m,5H), 7.69(s,1H)

Elemental analysis ( $\text{C}_{16}\text{H}_{22}\text{N}_2\text{OS HNO}_3$ )				
Calcd.(%):	C,54.37:	H,6.56:	N,11.89:	S,9.07
Found (%):	C,54.20:	H,6.54:	N,11.78:	S,8.92

I-9:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: (free base) 0.95(t,J=7.2Hz,3H), 1.30-1.80(m,2H), 2.69(t,J=7.4Hz,2H), 4.39(s,2H), 5.32(s,2H), 6.97-7.32(m,10H), 7.89(s,1H)

Elemental analysis ( $\text{C}_{20}\text{H}_{22}\text{N}_2\text{OS HNO}_3$ )				
Calcd.(%):	C,59.83:	H,5.77:	N,10.47:	S,7.99
Found (%):	C,59.66:	H,5.72:	N,10.32:	S,7.87

I-10:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: 4.54(s,2H), 5.26(s,2H), 7.18-7.42 (m,15H)

I-11:

 $^1\text{H-NMR}$  ( $\text{CDCl}_3$ -TMS) $\delta$ ppm: (free base) 1.02(t,J=7.0Hz,3H), 2.32(s,3H), 3.41(q,J=7.0Hz,2H), 5.29(s,2H), 6.99-7.26(m,5H)

Elemental analysis ( $C_{13}H_{13}BrN_2OS \cdot HNO_3$ )

Calcd. (%): C, 40.01; H, 4.13; Br, 20.48; N, 10.77; S, 8.22

Found (%): C, 39.86; H, 4.21; Br, 20.61; N, 10.61; S, 8.19

I-12:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: (free base) 1.23(t, J=7.5Hz, 3H), 2.69(q, J=7.5Hz, 2H), 4.47(s, 2H), 5.36(s, 2H), 6.99-7.31(m, 10H)

Elemental analysis ( $C_{19}H_{19}BrN_2OS$ )					
Calcd. (%):	C, 56.58:	H, 4.75:	Br, 19.81:	N, 6.95:	S, 7.95
Found (%):	C, 56.66:	H, 4.77:	Br, 19.62:	N, 7.04:	S, 7.84

I-13:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: (free base) 0.92(t, 7.2Hz, 3H), 1.67(m, 2H), 2.63(m, 2H), 4.56(s, 2H), 5.35(s, 2H), 7.00-7.30 (m, 10H)

Elemental analysis ( $C_{20}H_{21}BrN_2OS \cdot HCl$ )						
Calcd. (%):	C, 52.93:	H, 4.89:	Br, 17.61:	Cl, 7.81:	N, 6.17:	S, 7.07
Found (%):	C, 52.73:	H, 4.88:	Br, 17.57:	Cl, 7.75:	N, 6.14:	S, 7.14

I-14:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: 1.24(t, J=7.4Hz, 3H), 2.20(s, 6H), 2.52(s, 3H), 2.68(q, J=7.4Hz, 2H), 4.37(s, 2H), 5.30(s, 2H), 6.60-6.74(m, 3H), 7.10-7.30(m, 5H)

I-15:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: 1.06(t, J=7.0Hz, 3H), 1.24 (t, J=7.4Hz, 2H), 2.22(s, 6H), 2.53(s, 3H), 2.67(q, J=7.4Hz, 2H), 3.36(q, J=7.0Hz, 2H), 5.23(s, 2H)

I-16:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: 1.27(d, J=7.2Hz, 6H), 2.20(s, 6H), 2.53(s, 3H), 3.18(sept, 1H), 4.38(s, 2H), 5.30(s, 2H), 6.60-6.74(m, 3H), 7.12-7.31(m, 5H)

Elemental analysis ( $C_{23}H_{28}N_2OS \cdot 2/3H_2O$ )				
Calcd. (%):	C, 70.37:	H, 7.53:	N, 7.13:	S, 8.17
Found (%):	C, 70.34:	H, 7.31:	N, 7.21:	S, 7.89

I-17:


$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: 1.26(d, J=6.8Hz, 6H), 2.52(s, 3H), 3.17(sept, 1H), 4.37(s, 2H), 5.29(s, 2H), 6.96-7.30(m, 10H)

Elemental analysis ( $C_{21}H_{24}N_2OS$ )				
Calcd. (%):	C, 71.75:	H, 6.86:	N, 7.95:	S, 9.10
Found (%):	C, 71.47:	H, 6.88:	N, 7.84:	S, 8.86

I-18:

$^1H$ -NMR( $CDCl_3$ -TMS)  $\delta$ ppm: 1.03(t, J=7.0Hz, 3H), 1.25 (d, J=7.0Hz, 6H), 2.53(s, 3H), 3.17(sept, 1H), 3.35(q, J=7.0Hz, 2H), 5.22(s, 2H), 6.95-7.27(m, 5H)


Elemental analysis (C <sub>16</sub> H <sub>22</sub> N <sub>2</sub> OS HCl 1/3H <sub>2</sub> O)					
Calcd.(%):	C,57.73:	H,7.17:	Cl,10.65:	N,8.42:	S,9.63
Found (%):	C,57.92:	H,7.13:	Cl,10.70:	N,8.62:	S,9.56

**Example 19****1-benzoyloxymethyl-2,4-diethyl-5-phenylthioimidazole (I-19)**

In a mixture of 5 ml of dry tetrahydrofuran and 2.5 ml of dry hexamethylphosphoric triamide is dissolved 150 mg of 1-benzoyloxymethyl-4-ethyl-5-phenylthioimidazole (I-4) (0.46 mmol) obtained in example 4, and the resulting solution is chilled at -78°C. To the solution is added dropwise 1.64 M n-butyl lithium solution in 0.49 ml of hexane (0.8 mmol). Five minutes later, 110 mg of ethyl iodide (0.8 mmol) is added to the mixture, which mixture is allowed to stand for 10 minutes, when it is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate and filtered. The filtrate is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:2) to give 50 mg of the product (I-19) as an oil. Yield, 31%.

**I-19:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.20(t,J=7.4Hz,3H), 1.37 (t,J=7.4Hz,3H), 2.71(q,J=7.4Hz,2H), 2.83(q,J=7.4Hz,2H), 4.36(s,2H), 5.31(s,2H), 6.96-7.29(m,10H)


**Example 20****1-Benzoyloxymethyl-2-bromo-4-methyl-5-phenylthioimidazole (I-20)**

In 5 ml of dry tetrahydrofuran is dissolved 300 mg of 1-benzoyloxymethyl-4-methyl-5-phenylthioimidazole (I-3) (1 mmol) obtained in example 3, and the resulting solution is chilled at -78°C. To the solution is added dropwise 1.64 M solution of n-butyl lithium in 0.71 ml of hexane (1.16 mmol) and the mixture allowed to stand for 10 minutes, when 185 mg of bromine (1.16 mmol) is added thereto. Fifteen minutes later, the mixture is poured onto ice water. The mixture is neutralized with sodium bicarbonate and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The crude product is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:2) to give 360 mg of the product (I-20; mp. 55°C). Yield, 96%.

**I-20:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.33(s,3H), 4.45(s,2H), 5.36(s,2H), 6.99-7.30(m,10H)

Elemental analysis (C <sub>18</sub> H <sub>17</sub> BrN <sub>2</sub> SO)					
Calcd.(%):	C,55.53:	H,4.40:	Br,20.53:	N,7.20:	S,8.24
Found (%):	C,55.56:	H,4.53:	Br,20.26:	N,7.37:	S,8.31

**Example 21****1-Benzylloxymethyl-2-formyl-4-methyl-5-phenylthioimidazole hydrochloride (I-21)**

In 5 ml of dry tetrahydrofuran is dissolved 300 mg of 1-benzylloxymethyl-4-methyl-5-phenylthioimidazole (I-3) (1 mmol), and the solution is chilled at -78°C. To the solution is added dropwise 1.64 M solution of n-butyl lithium in 0.71 ml of hexane (1.16 mmol) and the mixture allowed to stand for 10 minutes, when 85 mg of dimethylformamide (1.16 mmol) is added thereto. Fifteen minutes later, the mixture is poured onto ice water. The mixture is extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate and concentrated. The crude product is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:2) to give 194 mg of the product (I-21) as an oil. Yield, 59%. This product is treated with hydrochloric acid/ethyl ether to give the hydrochloride as crystals melting at 97 - 98 °C.


**I-21:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.35(s,3H), 4.53(s,2H), 5.88(s,2H), 7.02-7.29(m,10H), 9.81(s,1H), IR(film): 1691, 1454, 1112, 1086cm<sup>-1</sup>

**Elemental analysis (C<sub>19</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>S HCl H<sub>2</sub>O)**

Calcd. (%) : C, 58.08 : H, 5.39 : Cl, 9.02 : N, 7.13 : S, 8.16

Calcd. (%) : C, 57.85 : H, 5.46 : Cl, 8.66 : N, 7.09 : S, 7.97


**Example 22****1-Benzylloxymethyl-2-ethoxycarbonyl-4-methyl-5-phenylthioimidazole nitrate (I-22)**

In 10 ml of dry tetrahydrofuran is dissolved 300 mg of 1-benzylloxymethyl-4-methyl-5-phenylthioimidazole (I-3) (1 mmol) obtained in Example 3, and the solution is chilled at -78°C. To the solution is added dropwise 1.64 M solution of n-butyl lithium in 0.88 ml of hexane (1.44 mmol) and the mixture allowed to stand for 10 minutes, when 157 mg of ethyl chloroformate (1.44 mmol) is added thereto. Thirty minutes later, the mixture is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate and filtered. The filtrate is concentrated, and the crude product is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1). The product is treated with nitric acid/ethyl ether to give 71 mg of the product (I-22) as crystals melting at 90 - 91°C. Yield, 17%.

**I-22:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: (free base) 1.45(t, J=7.0Hz, 3H), 2.36(s, 3H), 4.45(q, J=7.0Hz, 2H), 4.51(s, 2H), 5.90(s, 2H), 6.97-7.28(m, 10H), IR(film): 1712, 1474, 1237 cm<sup>-1</sup>


Elemental analysis (C <sub>21</sub> H <sub>22</sub> N <sub>2</sub> O <sub>3</sub> S HNO <sub>3</sub> )				
Calcd. (%):	C, 56.62:	H, 5.20:	N, 9.43:	S, 7.20
Found (%):	C, 56.79:	H, 5.29:	N, 9.42:	S, 7.28

**Example 23****2,5-Bis-phenylthio-1-benzyloxymethyl-4-methylimidazole (I-23)**

In 6 ml of dry tetrahydrofuran is dissolved 200 mg of 1-benzyloxymethyl-4-methyl-5-phenylthioimidazole (I-3) (0.64 mmol), and the solution is chilled at -78°C. To the solution is added dropwise 1.64 M solution of n-butyl lithium in 0.6 ml of hexane (0.98 mmol) and the mixture allowed to stand for 10 minutes, when 211 mg of diphenyl disulfide (9.7 mmol) is added thereto. Thirty minutes later, the mixture is poured onto ice water. The mixture is extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The crude product is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:2) to give 250 mg of the product (I-23) as an oil. Yield, 93%.

**I-23:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.38(s, 3H), 4.38(s, 2H), 5.46(s, 2H), 6.98-7.31(m, 15H)


**Example 24****1-Benzyloxymethyl-2-hydroxymethyl-4-methyl-5-phenylthioimidazole (I-24)**

To a solution of 130 mg of 1-benzyloxymethyl-2-formyl-4-methyl-5-phenylthioimidazole (I-21) (0.38 mmol), obtained in Example 21, in 5 ml of methyl alcohol is added 50 mg of sodium borohydride (1.3 mmol) under ice cooling and the mixture allowed to stand for 10 minutes, when acetic acid is added thereto. The mixture is neutralized with sodium bicarbonate and extracted with methylene chloride. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The residue is chromatographed on a column of silica gel, eluting with 2% methanol/ethyl acetate to give 124 mg of the product (I-24). Yield, 94%; mp. 87 - 88°C.

**I-24:**


<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.32(s, 3H), 3.00(brs., 1H), 4.37(s, 2H), 4.79(s, 2H), 5.44(s, 2H), 6.97-7.30(m, 10H), IR(film) 3172, 1475, 1454, 1438, 1074 cm<sup>-1</sup>

Elemental analysis (C <sub>19</sub> H <sub>20</sub> N <sub>2</sub> O <sub>2</sub> S)				
Calcd. (%):	C, 67.03:	H, 5.92:	N, 8.22:	S, 9.42
Found (%):	C, 66.83:	H, 6.05:	N, 8.31:	S, 9.33

**Example 25****1,2-Bis-benzyloxymethyl-4-methyl-5-phenylthioimidazole (I-25)**

To a solution of 20 mg of 1-benzyloxymethyl-2-hydroxymethyl-4-methyl-5-phenylthioimidazole (I-24) (0.06 mmol), obtained in Example 24, in 1 ml of dry dimethylformamide is added 4 mg of 60% oil suspension of sodium hydride (0.1 mmol) under ice cooling. Five minutes later, 15 mg of benzyl bromide (0.09 mmol) is added to the mixture, which is stirred at room temperature for 15 minutes. The mixture is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1) to give 25 mg of the product (I-25) as an oil. Yield, 95%.

I-25:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.37(s,2H), 4.34(s,2H), 4.58(s,2H), 4.75(s,2H), 5.47(s,2H), 7.00-7.33(m,15H)**Example 26****1-Benzyloxymethyl-2-carbamoyl-4-methyl-5-phenylthioimidazole (I-26)**


To a mixture of 10 ml of liquid ammonia and 1 ml of water in a sealing tube is added 200 mg of 1-benzyloxymethyl-2-ethoxycarbonyl-4-methyl-5-phenylthioimidazole (I-22) (0.5 mmol) obtained in Example 22 and the mixture is heated at 70°C for 3 hours. After evaporating the ammonia, the residue is chromatographed on a column of silica gel, eluting with ethyl acetate and n-hexane (1:1) to give 111 mg of desired product (I-26). Yield, 60%; mp. 153 - 154°C.

I-26:


<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.30(s,3H), 4.56(s,2H), 5.49(brs.,1H), 6.00(s,2H), 6.99-7.30(m,11H), IR(Nujol) 3338, 3208, 1657cm<sup>-1</sup>



Elemental analysis (C <sub>19</sub> H <sub>19</sub> N <sub>3</sub> O <sub>2</sub> S)				
Calcd.(%):	C,64.57:	H,5.42:	N,11.89:	S,9.07
Found (%):	C,64.51:	H,5.49:	N,11.89:	S,9.04

**Example 27****1-Benzyloxymethyl-2,4-dibromo-5-phenylthioimidazole (I-27)**

To a solution of 50 mg of 1-benzyloxymethyl-5-phenylthioimidazole nitrate (I-5) (0.14 mmol), obtained in Example 5, in 2 ml of acetic acid is added 70 mg of bromine (0.43 mmol), and the mixture is stirred at room temperature for 1 hour and poured onto ice water. The resultant mixture is neutralized with sodium bicarbonate and extracted with methylene chloride. The extract is washed with water, dried over sodium sulfate, filtered and the filtrate is concentrated. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (3:1) to give 40 mg of the product (I-27). Yield, 64%; mp. 89-90°C.

**I-27:**<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 4.45(s,2H), 5.43(s,2H), 7.00-7.30 (m,10H)**Example 28****1-Benzyloxymethyl-2-hydroxy-4-methyl-5-phenylthioimidazole (I-28)**


To a mixture of 15 mg of 1-benzyloxymethyl-2-bromo-4-methyl-5-phenylthioimidazole (I-20) (0.04 mmol), obtained in example 20, and 3 ml of 2-methoxyethyl ether is added 1 ml of 1 M sodium methoxide solution in methanol, and the resultant mixture is heated at 130°C for 5 hours. After evaporating the 2-methoxyethyl ether, the residue is chromatographed on a column of silica gel, eluting with 3% methanol/methylene chloride. The crude product is recrystallized from ethyl acetate/isopropyl ether to give 9 mg of the product (I-28). Yield, 71%; mp. 147 - 148°C.

**I-28:**<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.22(s,2H), 4.54(s,2H), 5.11(s,2H), 7.11-7.26(m,10H), 9.94(br,1H), IR(Nujol) 3100, 1686, 1457, 1369cm<sup>-1</sup>

Elemental analysis (C <sub>18</sub> H <sub>18</sub> N <sub>2</sub> O <sub>2</sub> S)				
Calcd.(%):	C,66.23:	H,5.56:	N,8.58:	S,9.82
Found (%):	C,66.11:	H,5.68:	N,8.65:	S,9.55

## Example 29

## 5-(3,5-Dimethylphenylthio)-4-ethyl-1-[(2-hydroxyethoxy)-methyl]-2-methylimidazole (I-29)



(1) To a solution of 500 mg of 5-(3,5-dimethylphenylthio)-4-ethyl-2-methylimidazole (2 mmol) in 5 ml of dry dimethylformamide is added 120 mg of 60% oil suspension of sodium hydride (3 mmol) under ice cooling and allowed to stand for 5 minutes. After the addition of 820 mg of iodomethyl trimethylsilyloxyethyl ether (3 mmol) [Tetrahedron Letters, No. 35, pp. 3263 - 3264, 1979], the mixture is allowed to stand for 15 minutes, when it is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and concentrated in vacuo. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (2:1) to give 170 mg of 5-(3,5-dimethylphenylthio)-4-ethyl-2-methyl-1-trimethylsilyloxyethoxymethyl-imidazole 4 as an oil. Yield, 21%.

4:  
<sup>1</sup>H-NMR(CDCl<sub>3</sub>)δppm: 0.00(s,9H), 1.27(t,J=7.6Hz,3H), 2.15(s,6H), 2.67(s,3H), 2.74(q,J=7.6Hz,2H), 3.28(m,2H), 3.47(m,2H), 5.31(s,2H), 6.54(s,2H), 6.74(s,1H)

(2) Preparation of I-29:

To a mixture of 0.85 ml of 10% aqueous solution of potassium fluoride, 0.85 ml of 10% aqueous solution of potassium bicarbonate and 3.4 ml of dimethylformamide is added 170 mg of 5-(3,5-dimethylphenylthio)-4-ethyl-2-methyl-1-trimethylsilyloxyethoxymethyl-imidazole 4 obtained in (1) above, and the resultant mixture is stirred at room temperature for 30 minutes. The mixture is combined with water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and concentrated. The residue is chromatographed on a column of silica gel, eluting with 5% methanol/ethyl acetate. The crude product is recrystallized from n-hexane to give 80 mg of the product (I-29) as crystals melting at 87 - 89°C. Yield, 58%.


I-29:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.24(t,J=7.4Hz,3H), 2.22(s,6H), 2.54(s,3H), 2.69(q,J=7.4Hz,2H), 5.29(s,2H), 6.61(s,2H), 6.76(s,1H)

Elemental analysis(C<sub>17</sub>H<sub>24</sub>N<sub>2</sub>O<sub>2</sub>S)

Calcd. (%):C, 63.72;H, 7.55;N, 8.74;S, 10.01



Found (%):C, 63.68;H, 7.56;N, 8.81;S, 9.81

**Example 30****5-(2,4-Difluorophenylthio)-4-isopropyl-1-benzyloxymethyl-2-methylimidazole (I-30)**

The reaction is effected in the same manner as in examples 1 and 2 to give the product (I-30).

**I-30:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.23(d,J=7.2Hz,6H), 2.50(s,3H), 3.18(sept,1H), 4.42(s,2H), 5.32(s,2H), 6.70-6.80(m,3H), 7.14.28(m,5H)

**Example 31****5-(2,4-Difluorophenylthio)-1-ethoxymethyl-4-isopropyl-2-methylimidazole (I-31) and 4-(2,4-difluorophenylthio)-1-ethoxymethyl-5-isopropyl-2-methylimidazole (I-86)**

To a solution of 300 mg of 5-(2,4-difluorophenylthio)-4-isopropyl-2-methylimidazole (1.2 mmol) in 3 ml of dry dichloromethane is added 240 mg of triethylamine (2.4 mmol). To the mixture is added dropwise 205 mg of chloromethylethyl ether (2.4 mmol) under ice cooling. After 1-hour-reaction at room temperature, to the reaction mixture is added ice water, which is followed by the addition of aqueous sodium hydrogencarbonate and extraction with methylene chloride. The extract is dried over sodium sulfate, filtered and concentrated in vacuo. The residue containing two products is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1) to separate each product.

The former eluate gives 5 mg of 4-(2,4-difluorophenylthio)-1-ethoxymethyl-5-isopropyl-2-methylimidazole (I-86). Yield, 2%.

The later eluate gives 310.0 mg of 5-(2,4-difluorophenylthio)-1-ethoxymethyl-4-isopropyl-2-methylimidazole (I-31). Yield, 84%.

**I-31:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.05(t,J=7.0Hz,3H), 1.22 (d,J=7.0Hz,6H), 2.52(s,3H), 3.17(sept,1H), 3.38(q,J=7.0Hz,2H), 5.24(s,2H), 6.70-6.90(m,3H)


**I-86:**

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.02(t,J=6.8Hz,3H), 1.22 (d,J=7.0Hz,6H), 2.52(s,3H), 3.13(sept,1H), 3.35(q,J=6.8Hz,2H), 5.21(s,2H), 6.50-7.50(m,3H)

**Examples 32 - 36**

In examples 32 to 36, the reaction is effected in the same manner as in Examples 1 - 2 to give the products I-32 to I-36 as summarized in Table 2 below.

Table 2



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
32	S	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 169-170°C (H <sub>2</sub> SO <sub>4</sub> salt)	1 - 2 2
33	S	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>3</sub>	-CH <sub>3</sub>	mp 163-164°C (H <sub>2</sub> SO <sub>4</sub> salt)	1 - 3 3
34	S	-C(CH <sub>3</sub> ) <sub>3</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 67-68°C	1 - 3 4
35	S	-C(CH <sub>3</sub> ) <sub>3</sub>		-CH <sub>3</sub>	-CH <sub>3</sub>	mp 74-75°C	1 - 3 5
36	S	-C(CH <sub>3</sub> ) <sub>3</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 3 6

Physicochemical properties of the compounds as described in Table 2 are shown below.

1-32:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.26(m, J=7Hz, 6H), 1.65(t, J=7Hz, 3H), 2.12(s, 6H), 2.53(s, 3H), 3.17(m, J=7Hz, 1H), 3.36(q, J=7Hz, 2H), 5.22(s, 2H), 6.59-6.74(m, 3H)

Elemental analysis (C<sub>18</sub>H<sub>26</sub>N<sub>2</sub>OS H<sub>2</sub>SO<sub>4</sub>)

Calcd.:	C, 51.90:	H, 6.77:	N, 6.73:	S, 15.40
Found:	C, 51.78:	H, 6.78:	N, 6.71:	S, 15.66

I-33:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.26(m, J=7Hz, 6H), 2.22(s, 6H), 2.52 (s, 3H), 3.17(m, J=7Hz, 1H), 3.18(s, 3H), 5.18(s, 2H), 6.59-6.74(m, 3H)Elemental analysis (C<sub>17</sub>H<sub>24</sub>H<sub>2</sub>OS H<sub>2</sub>SO<sub>4</sub>)

Found :	C, 50.72:	H, 6.51:	N, 6.96:	S, 15.93
Calcd.:	C, 50.84:	H, 6.62:	N, 6.81:	S, 15.76

I-34:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.05(t, J=7Hz, 3H), 1.39(s, 9H), 2.21(s, 6H), 3.51(s, 3H), 3.38 (q, J=7Hz, 2H), 5.17(s, 2H), 6.53-6.73(m, 3H)Elemental analysis (C<sub>18</sub>H<sub>28</sub>N<sub>2</sub>OS H<sub>2</sub>SO<sub>4</sub>)

Calcd.:	C, 53.00:	H, 7.02:	N, 6.51:	S, 14.89
Found :	C, 53.14:	H, 7.05:	N, 6.61:	S, 14.98

I-35:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.39(s, 9H), 2.21(s, 6H), 2.51(s, 3H), 3.20(s, 3H), 5.13(s, 2H), 6.53-6.73(m, 3H)Elemental analysis (C<sub>18</sub>H<sub>28</sub>N<sub>2</sub>OS H<sub>2</sub>SO<sub>4</sub>)


Calcd.:	C, 51.80:	H, 6.77:	N, 6.73:	S, 15.40
Found :	C, 51.66:	H, 6.88:	N, 6.66:	S, 15.33

I-36:


<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.39(s, 9H), 2.50(s, 3H), 4.39(s, 2H), 5.24(s, 2H), 6.90-7.30(m, 10H)

## Example 37

1-Benzoyloxymethyl-4-isopropyl-2-methyl-5-phenylselenenylimidazole (I-37) and 1-benzoyloxymethyl-4-phenylselenenyl-2-methyl-5-isopropyl-imidazole (I-87)



I - 37



I - 87

## (1) 2-methyl-4-isopropyl-5-phenylselenenyl-imidazole and 2-methyl-4-phenylselenenyl-5-isopropyl-imidazole

To a solution of 960 mg of benzeneselenenol (6.0 mmol) in 10 ml of dry dimethylformamide is added 400 mg of 60% oil suspension of sodium hydride (10 mmol) under ice cooling and the mixture allowed to stand for 10 minutes. After the addition of 1 g of 2-methyl-4-isopropyl-5-iodoimidazole (4 mmol), the mixture is heated at 140°C in a nitrogen atmosphere for 1 hour. The resultant mixture is mixed with ice water and dry ice and extracted with diethyl ether. The extract is washed with water, dried over sodium sulfate and filtered. The filtrate is concentrated in vacuo, and the crude product is chromatographed on a column of silica gel, eluting with 3% methanol/methylene chloride. The product is recrystallized from isopropyl ether to give 900 mg of the desired 2-methyl-4-isopropyl-5-phenylselenenyl-imidazole and 2-methyl-4-phenylselenenyl-5-isopropyl-imidazole. Yield 80%; mp. 146 - 147°C.

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.24(d, J=7.20Hz, 6H), 2.41(s, 3H), 3.21(sept, 1H), 7.18(m, 5H)

(2) 1-benzyloxymethyl-4-isopropyl-2-methyl-5-phenylselenenyl-imidazole (I-37) and 1-benzyloxymethyl-2-methyl-4-phenylselenenyl-5-isopropyl-imidazole (I-87)

To a solution of 400 mg of 2-methyl-4-isopropyl-5-phenylselenenyl-imidazole (1.4 mmol) in 4 ml of dry dimethylformamide is added 70 mg of 60% oil suspension of sodium hydride (1.8 mmol) under ice cooling and the mixture is allowed to stand for 5 minutes followed by the addition of 270 mg of benzyloxymethyl chloride (1.7 mmol). The mixture is stirred for 30 minutes under ice cooling, mixed with ice water and extracted with diethyl ether. The extract is washed with water, dried over sodium sulfate and filtered. The filtrate is concentrated in vacuo. The crude product is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1). The product is recrystallized from n-hexane to give 297 mg of the objective 2-methyl-4-phenylselenenyl-5-isopropyl-imidazole (I-87) and 1-benzyloxymethyl-4-isopropyl-2-methyl-5-phenylselenenyl-imidazole (I-37). Yield, 52%; mp. 94 - 96°C


I-37:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.27(d,J=7.0Hz,6H), 2.55(s,3H), 3.21(sept,1H), 4.36(s,3H), 5.36(s,2H), 7.10-7.30(m,10H)

Elemental analysis (C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> OSe)			
Calcd.(%):	C,63.15:	H,6.06:	N,7.01
Found (%):	C,63.12:	H,6.04:	N,6.97

I-87:


<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.34(t,J=7.0HZ,6H), 2.53(s,3H), 3.15(sept,1H), 4.54(s,2H), 5.29(s,2H), 7.20-7.50(m,10H)

Example 381-Ethoxymethyl-4-isopropyl-2-methyl-5-phenylselenenyl-imidazole (I-38)

The titled compound (I-38) was prepared as an oil in the same manner as that described in Example 37.

I-38:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.03(t,J=7.0HZ,3H), 1.27 (d,J=7.0HZ,6H), 3.21(sept,1H), 3.34(q,2H), 5.28(s,2H), 7.10-7.26(m,5H)

Example 391-(2-Hydroxyethoxymethyl)-4-isopropyl-2-methyl-5-phenylselenenyl-imidazole (I-39)

Compound (I-39) was prepared according to the synthetic route B.

To a solution of 560 mg of 2-methyl-4-isopropyl-5-phenylselenenyl-imidazole (2 mmol) in 5.6 ml of dry dimethylformamide is added 120 mg of 60% oil suspension of sodium hydride (3 mmol) under ice cooling and the mixture is allowed to stand for 5 minutes. To the mixture is added 820 mg of iodomethyl trimethylsilyloxyethyl ether (3 mmol) and allowed to stand for 10 minutes, when the reaction mixture is poured onto ice water and extracted with ethyl ether. The extract is washed with water, dried over sodium sulfate, filtered and concen-


trated in vacuo. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (2:1) to give 70 mg of 1-(2-hydroxyethoxymethyl)-4-isopropyl-2-methyl-5-phenylselenenyl-imidazole as an oil. Yield, 10%.

I-39:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.27(d, J=6.8Hz, 6H), 2.56(s, 3H), 2.97(t, J=6.6Hz, 2H), 3.21(sept, 1H), 3.52(t, J=6.6Hz, 2H), 5.33(s, 2H), 6.90.26(m, 5H)

#### Example 40

##### 1-Benzylloxymethyl-4-isopropyl-2-methyl-5-phenylsulfinyl-imidazole (I-40)



To a solution of 500 mg of 1-benzylloxymethyl-4-isopropyl-2-methyl-5-phenylthio-imidazole (1.4 mmol) in 10 ml of methylene chloride is added 370 mg of m-chloroperbenzoic acid (2.1 mmol) under ice cooling, and the resultant mixture is stirred at room temperature for 3 hours. The mixture is mixed with aqueous sodium thiosulfate and aqueous sodium hydrogencarbonate and extracted with methylene chloride. The extract is washed with water, dried over sodium sulfate, filtered and concentrated in vacuo. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1) to give 471 mg of the objective 1-benzylloxymethyl-4-isopropyl-2-methyl-5-phenylsulfinyl-imidazole as crystals melting at mp. 72 - 74°C. Yield, 54%.


I-40:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.36(d, d, J=7.2Hz, 7.0Hz, 6H), 2.38(s, 3H), 3.33(sept, 1H), 4.12(d, J=11.6Hz, 1H), 4.21(d, J=11.6Hz, 1H), 5.18(s, 2H), 7.09-7.52(m, 10H)

Elemental analysis (C <sub>21</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub> S)				
Calcd. (%):	C, 68.45:	H, 6.57:	N, 7.60:	S, 8.70
Found (%):	C, 68.31:	H, 6.65:	N, 7.65:	S, 8.47

#### Example 41

##### 1-Benzylloxymethyl-4-isopropyl-2-methyl-5-phenylsulfonyl-imidazole (I-41)




To a solution of 1 g of 1-benzylloxymethyl-4-isopropyl-2-methyl-5-phenylthio-imidazole (2.8 mmol) in 40 ml of methylene chloride is added 1.84 g of m-chloroperbenzoic acid (10.7 mmol) under ice cooling, and the resultant mixture is stirred at room temperature. Six hours later, the reaction mixture is mixed with aqueous sodium thiosulfate and aqueous sodium hydrogencarbonate and extracted with methylene chloride. The extract is washed with water, dried over sodium sulfate, filtered and concentrated in vacuo. The residue is chromatographed on a column of silica gel, eluting with ethyl acetate/n-hexane (1:1) to give 580 mg of the objective 1-benzylloxymethyl-4-isopropyl-2-methyl-5-phenylsulfonyl-imidazole as crystals melting at mp. 116 - 117°C. Yield, 53%.

I-41:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.26(d, J=7.2Hz, 6H), 2.42(s, 3H), 3.76(sept, 1H), 4.41(s, 2H), 5.59(s, 2H), 7.17-7.58(m, 8H), 7.87-7.91(m, 2H)

Elemental analysis (C<sub>21</sub>H<sub>24</sub>N<sub>2</sub>O<sub>3</sub>S H<sub>2</sub>O)

Calcd.(%):	C, 62.66:	H, 6.51:	N, 6.96:	S, 7.98
Found (%):	C, 62.73:	H, 6.25:	N, 7.01:	S, 7.83

Example 425-Benzyl-1-benzoyloxymethyl-4-isopropyl-2-methylimidazole (I-42)(1) N-acetylphenylalanine methyl ester 4

To a solution of 2.73 g of L-phenylalanine methyl ester hydrochloride (12.7 mmol) in 5.00 g of pyridine (63.2 mmol) is dropwise added 12.90 g of acetic anhydride (126.0 mmol) under ice cooling. The mixture is allowed to stand for 2 hours and concentrated in vacuo. The residue is extracted with ethyl acetate. The extract is washed with 5% hydrochloric acid, saturated aqueous sodium bicarbonate and saturated brine in order, dried over sodium sulfate, filtered and concentrated in vacuo to give 2.34 g of the objective N-acetylphenylalanine methyl ester 4. Yield, 83.6%.

I-42:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.99(s, 3H), 3.13(m, 2H), 3.73(s, 3H), 4.89(m, 1H), 5.89(br, 1H), 7.06-7.33(m, 5H)

Elemental analysis (C<sub>12</sub>H<sub>15</sub>NO<sub>3</sub>)

Calcd.(%);	C, 65.14:	H, 6.83:	N, 6.33
Found (%);	C, 65.12:	H, 6.80:	N, 6.31

(2) 4-Acetylamino-2-methyl-5-phenyl-3-pentanone 5

A solution of 2.30 g of N-acetylphenylalanine methyl ester (10.40 mmol) in 15 ml of tetrahydrofuran is chilled at -60°C with dry ice/acetone bath. To the mixture is added dropwise 10.40 ml of 2 M solution of isopropylmagnesium chloride in tetrahydrofuran (20.8 mmol) with stirring and the mixture allowed to stand for 15 minutes and returned to room temperature. The mixture was stirred for 3 hours. To the mixture is added saturated aqueous ammonium chloride under ice cooling, followed by the addition of 5% hydrochloric acid and the mixture extracted with ethyl acetate. The extract is washed with saturated brine, dried over sodium sulfate, filtered and the filtrate concentrated in vacuo. The resulting crude product is chromatographed on a column of silica gel, eluting with methylene chloride and ethyl acetate (4:1) to give 1.40 g of the objective 4-acetylamino-2-methyl-5-phenyl-3-pentanone 5. Yield, 57.7%.



5:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.00(d,J=6.6Hz,3H),

1.09(d,J=7.2Hz,3H), 1.97(s,3H), 2.68(m,1H), 3.04(m,2H), 5.06(m,1H), 6.07(br,1H), 7.08-7.34(m,5H)

Elemental analysis (C<sub>14</sub>H<sub>18</sub>NO<sub>2</sub> 0.1H<sub>2</sub>O)

Calcd.(%);	C,71.52;	H,8.23;	N,5.96
Found (%);	C,71.62;	H,8.08;	N,6.03

(3) 5-Benzyl-4-isopropyl-2-methylimidazole 6

To a solution of 1.00 g of 4-acetylamino-2-methyl-5-phenyl-3-pentanone (4.29 mmol) in 20 ml of acetic acid is added 5.00 g of ammonium acetate (64.9 mmol), and the resultant mixture is heated under reflux. One hour later, the reaction mixture is concentrated in vacuo, neutralized with aqueous sodium bicarbonate and extracted with ethyl acetate. The extract is washed with saturated brine, dried over sodium sulfate, filtered and the filtrate is concentrated. The crude product is recrystallized from ether/n-hexane to give 0.90 mg of the objective 5-benzyl-4-isopropyl-2-methylimidazole 6. Yield, 97.5%.

6:<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.23(d,J=7.0Hz,6H), 2.29(m,1H), 3.86(s,2H), 7.13-7.31(m,5H)Elemental analysis (C<sub>14</sub>H<sub>18</sub>N<sub>2</sub>)

Calcd.(%);	C,78.46;	H,8.47;	N,13.07
Found (%);	C,78.20;	H,8.50;	N,12.89

## (4) 5-Benzyl-1-benzyloxymethyl-4-isopropyl-2-methylimidazole (I-42)

To a solution of 980 mg of 5-benzyl-4-isopropyl-2-methylimidazole (4.58 mmol) in 9.8 ml of dry dimethylformamide is added 238 mg of 60% oil suspension of sodium hydride (5.95 mmol) under ice cooling. Ten minutes later, 935 mg of benzyloxymethyl chloride (5.97 mmol) is dropwise added. One hour later, the reaction mixture is poured onto ice water and extracted with ethyl acetate. The extract is washed with saturated brine, dried over sodium sulfate, filtered and the filtrate is concentrated in vacuo. The crude product is chromatographed on a column of silica gel, eluting with ethyl acetate and n-hexane (1:1) to give 820 mg of the objective 5-benzyl-1-benzyloxymethyl-4-isopropyl-2-methylimidazole (I-42). Yield, 54.17%.

I-42:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.28(d,J=6.6Hz,6H), 2.38(s,3H), 2.93(m,1H), 3.96(s,2H), 4.32(s,2H), 4.96(s,2H), 7.01-7.39(m,10H)Elemental analysis (C<sub>22</sub>H<sub>26</sub>N<sub>2</sub>O 0.8H<sub>2</sub>O)


Calcd.(%);	C,75.74;	H,7.97;	N,8.03
Found (%);	C,75.78;	H,7.75;	N,8.00

## Examples 43 - 70

In Examples 43 to 70, desired compounds (I-43) - (I-70) and (I-88) - (I-95) as shown in the following Tables 3 - 8 were prepared according to the synthetic route D in the same manner as described in Example 42 above. Among them, compounds (I-88) - (I-95) are position isomers of compounds (I-61), (I-62) and (I-67) - (I-72), respectively. The isomers are separated from each other at an appropriate reaction step of each Example, said step corresponding to Example 42, (2) (i.e., step 2 of synthetic route D). Example 61: compounds I-61 and I-


88; Example 62: compounds I-62 and I-89; Example 67: compounds I-67 and I-90; Example 68: compounds I-68 and I-91; Example 69: compounds I-69 and I-92; Example 70: compounds I-70 and I-93; Example 71: compounds I-71 and I-94; and Example 72: compounds I-72 and I-95.

Table 3




Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
43	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub> CH <sub>2</sub> OAc	-CH <sub>3</sub>	oil	1 - 4 3
44	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 60-61°	1 - 4 4
45	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub> CH <sub>2</sub> OH	-CH <sub>3</sub>	mp 130-131°	1 - 4 5
46	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 4 6
47	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 4 7
48	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 55-56°	1 - 4 8
49	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 4 9

Table 4




Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
50	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1-50
51	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1-51
52	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 225-226°C	1-52
53	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	mp 203-204°C	1-53
54	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>3</sub>	oil	1-54

Table 5




Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
55	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>3</sub>	-CH <sub>3</sub>	oil	1 - 55
56	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	-CH <sub>3</sub>	oil	1 - 56
57	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 57
58	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 58
59	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 59
60	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 60
61	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	mp 104-105°C	1 - 61

Table 6




Example No.	X	$R^1$	$R^2$	$R^3$	$R^4$	mp(°C)	Compound No.
62	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 80-81°C	1 - 6 2
63	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub>	-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 6 3
64	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	-CH <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 6 4
65	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 6 5
66	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 6 6
67	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 40-41°C	1 - 6 7

Table 7



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
68	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 68
69	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 69
70	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 70

Table 8



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
61	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 8 8
62	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	mp 80-81°C	1 - 8 9
67	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 9 0
68	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 9 1
69	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 9 2
70	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 9 3
71	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	oil	1 - 9 4
72	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1 - 9 5

Physicochemical properties of compounds shown in the Tables 3 to 8 above are shown below.

1-43:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.28(d, J=7.0Hz, 6H), 2.05(s, 3H), 2.42(s, 3H), 2.92(m, 1H), 3.39(m, 2H), 4.02(m, 4H), 4.97(s, 2H), 7.07-7.32(m, 5H)

Elemental analysis (C <sub>19</sub> H <sub>26</sub> N <sub>2</sub> O <sub>3</sub> 0.2H <sub>2</sub> O)			
Calcd.:	C,68.32:	H,7.97:	N,8.39
Found :	C,68.21:	H,7.97:	N,8.44

I-44:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.08(t,J=7.0Hz,3H),  
 1.29(d,J=7.0Hz,6H), 2.44(s,3H), 2.93(m,1H),  
 3.30(q,J=7.0Hz,2H), 3.99(s,2H), 4.92(s,2H), 7.07-7.32(m,5H)

Elemental analysis (C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> O 0.05H <sub>2</sub> O)			
Calcd.:	C,74.71:	H,8.89:	N,10.25
Found :	C,74.64:	H,8.82:	N,10.20

I-45:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.29(d,J=7.0Hz,6H), 2.44(s,3H), 2.93(m,1H), 3.34(m,2H), 3.57(m,2H), 3.99(s,2H),  
 5.00(s,2H), 7.08-7.32(m,5H)

Elemental analysis (C <sub>17</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub> )			
Calcd.:	C,70.80:	H,8.39:	N,9.71
Found :	C,70.71:	H,8.41:	N,9.67

I-46:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm:1.28(d,J=6.6Hz,6H), 2.38(s,3H), 2.91(m,1H), 3.76(s,3H), 3.89(s,2H), 4.33(s,2H),  
 4.96(s,2H), 6.75-7.35(m,9H)

Elemental analysis (C <sub>23</sub> H <sub>28</sub> N <sub>2</sub> O <sub>2</sub> 0.5H <sub>2</sub> O)			
Calcd.:	C,73.96:	H,7.83:	N,7.50
Found :	C,73.93:	H,7.76:	N,7.66

I-47:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.10(t,J=7.0Hz,3H),  
 1.28(d,J=6.8Hz,6H), 2.42(s,3H), 2.92(m,1H),  
 3.31(q,J=7.0Hz,2H), 3.78(s,3H), 3.92(s,2H), 4.91(s,2H), 6.79-7.03(m,4H)

Elemental analysis (C <sub>18</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub> 0.2H <sub>2</sub> O)			
Calcd.:	C,70.65:	H,8.70:	N,9.15
Found :	C,70.72:	H,8.55:	N,9.41

I-48:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.08(t,J=7.0Hz,3H),  
 1.28(d,J=7.0Hz,6H), 1.42(s,3H), 2.93(m,1H),  
 3.30(q,J=7.0Hz,2H), 3.92(s,2H), 4.91(s,2H), 5.03(s,2H), 6.86-7.45(m,9H)

Elemental analysis (C <sub>24</sub> H <sub>30</sub> N <sub>2</sub> O <sub>2</sub> 0.4DMF 0.2H <sub>2</sub> O)			
Calcd.:	C,73.58:	H,8.14:	N,8.17
Found :	C,73.73:	H,8.19:	N,7.89



I-49:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.28(d,J=6.8Hz,6H), 2.38(s,3H), 2.92(m,1H), 3.89(s,2H), 4.32(s,2H), 4.96(s,2H), 5.01(s,2H), 6.82-7.43(m,14H)

Elemental analysis (C <sub>29</sub> H <sub>32</sub> N <sub>2</sub> H <sub>2</sub> 0.5H <sub>2</sub> O)			
Calcd.:	C,77.47:	H,7.40:	N,6.23
Found :	C,77.51:	H,7.22:	N,6.40

I-50:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.29(d,J=6.6Hz,6H), 2.22(s,6H), 2.41(s,3H), 2.93(m,1H), 3.87(s,2H), 4.34(s,2H), 4.97(s,2H), 6.62(s,2H), 6.80(s,1H), 7.18-7.40(m,5H)

Elemental analysis (C <sub>24</sub> H <sub>30</sub> N <sub>2</sub> O 0.3H <sub>2</sub> O)			
Calcd.:	C,78.35:	H,8.38:	N,7.61
Found :	C,78.26:	H,8.38:	N,7.68

I-51:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.13(t,J=7.0Hz,3H), 1.32 (d,J=7.0Hz,6H), 2.28(s,6H), 2.47(s,3H), 2.95(m,1H), 3.35(q,J=7.0Hz,2H), 3.94(s,2H), 4.96(s,2H), 6.73(s,2H), 6.88(s,1H)

Elemental analysis (C <sub>18</sub> H <sub>28</sub> N <sub>2</sub> O 0.4H <sub>2</sub> O)			
Calcd.:	C,74.18:	H,9.44:	N,9.11
Found :	C,74.20:	H,9.40:	N,9.05

I-52:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.10(t,J=7.0Hz,3H), 1.29 (d,J=6.6Hz,6H), 2.44(s,3H), 2.95(m,1H), 3.31(q,J=7.0Hz,2H), 3.92(s,2H), 4.92(s,2H), 6.74-6.93(m,4H)

Elemental analysis (C <sub>17</sub> H <sub>24</sub> N <sub>2</sub> O <sub>2</sub> 0.1H <sub>2</sub> O)			
Calcd.:	C,70.36:	H,8.41:	N,9.65
Found.:	C,70.38:	N,8.37:	H,9.62

I-53:

<sup>1</sup>H-NMR(CD<sub>3</sub>OD-TMS)ppm: 1.22(d,J=6.6Hz,6H), 2.33(s,3H), 2.91(m,1H), 3.86(s,2H), 4.35(s,2H), 5.07(s,2H), 6.63-6.86(m,4H), 7.18-7.35(m,5H)

Elemental analysis (C <sub>22</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub> 0.6H <sub>2</sub> O)			
Calcd.:	C,73.14:	H,7.59:	N,7.75
Found :	C,73.23:	H,7.61:	N,7.48

I-54:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.05(d,J=6.2Hz,6H), 1.27 (d,J=7.0Hz,6H), 2.42(s,3H), 2.90(m,1H), 3.49(m,1H), 3.99(s,2H), 4.89(s,2H), 7.09-7.33(m,5H)

Elemental analysis (C <sub>18</sub> H <sub>26</sub> N <sub>2</sub> O 0.5H <sub>2</sub> O)			
Calcd.:	C73.18,	H9.21,	N9.48
Found.:	C73.31,	H9.06,	N9.59

I-55:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.28(d,J=7.0Hz,6H), 2.43(s,3H), 2.93(m,1H), 3.14(s,3H), 3.99(s,2H), 4.86(s,2H), 7.08-7.31(m,5H)

I-56:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 0.83(t,J=7.4Hz,3H), 1.28(d,J=7.2Hz,6H), 1.45(m,2H), 2.42(s,3H), 2.81(m,1H), 3.17(q,J=6.4Hz,2H), 3.99(s,2H), 4.91(s,2H), 7.08-7.33(m,5H)

Elemental analysis (C <sub>18</sub> H <sub>26</sub> N <sub>2</sub> O 0.3H <sub>2</sub> O)			
Calcd.:	C,74.09:	H,9.19:	N,9.60
Found.:	C,74.06:	H,9.20:	N,9.61

I-57:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.05(t,J=7.0Hz,3H), 1.27(d,J=7.2Hz,6H), 2.44(s,3H), 2.88(m,1H), 3.31(q,J=7.0Hz,2H), 3.99(s,2H), 4.92(s,2H), 7.04(d,J=6.2Hz,2H), 8.50(d,J=5.8Hz,2H)

I-58:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.06(t,J=7.0Hz,3H), 1.28(d,J=6.8Hz,6H), 2.44(s,3H), 2.90(m,1H), 3.31(q,J=7.0Hz,2H), 4.00(s,2H), 4.94(s,2H), 7.18-7.74(m,2H), 8.39-8.53(m,2H)

I-59:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.24(d,J=7.4Hz,6H), 2.36(s,3H), 3.05(m,1H), 3.95(s,2H), 4.51(s,2H), 5.22(s,2H), 7.15-7.60(m,7H), 8.39-8.52(m,2H)

I-60:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.05(t,J=7.0Hz,3H), 1.25(d,J=7.0Hz,6H), 2.42(s,3H), 2.93(m,1H), 3.33(q,J=7.0Hz,2H), 4.16(s,2H), 5.08(s,2H), 6.95-7.62(m,3H), 8.50-8.55(m,1H)

Elemental analysis (C <sub>18</sub> H <sub>23</sub> N <sub>3</sub> O 0.3H <sub>2</sub> O)			
Calcd.:	C,68.93:	H,8.53:	N,15.07
Found.:	C,68.83:	H,8.35:	N,15.05

I-61:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.02(d,J=7.0Hz,6H), 2.20(s,6H), 2.26(s,3H), 2.28(m,1H), 2.37(s,3H), 3.93(s,2H), 4.35(s,2H), 5.08(s,2H), 6.83(s,2H), 7.23-7.40(m,5H)

Elemental analysis (C <sub>25</sub> H <sub>32</sub> N <sub>2</sub> O 0.2H <sub>2</sub> O)			
Calcd.:	C,78.99:	H,8.59:	N,7.37
Found.:	C,78.52:	H,8.28:	N,7.17

I-62:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.02(d,J=6.8Hz,6H), 1.18(t,J=7.0Hz,3H), 2.21(s,6H), 2.21(m,1H), 2.27(s,3H), 2.45(s,3H), 3.37(q,J=7.0Hz,2H), 3.94(s,2H), 5.03(s,2H), 6.84(s,2H)

Elemental analysis (C <sub>20</sub> H <sub>30</sub> N <sub>2</sub> O 0.2H <sub>2</sub> O)			
Calcd.:	C,75.52:	H,9.63:	N,8.81
Found.:	C,75.45:	H,9.42:	N,8.71

I-63:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.19(d,J=7.0Hz,6H), 2.38(s,3H), 2.70-3.01(m,5H), 4.48(s,2H), 4.98(s,2H), 7.01-7.40(m,10H)

## EP 0 552 060 A1

Elemental analysis (C <sub>23</sub> H <sub>28</sub> N <sub>2</sub> O 0.6H <sub>2</sub> O)			
Calcd.:	C,76.89:	H,8.19:	N,7.80
Found :	C,76.74:	H,8.07:	N,7.80

## I-64:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.18(d,J=7.2Hz,6H), 1.20(t,J=7.0Hz,3H), 2.42(s,3H), 2.69-3.02(m,5H), 3.46(q,J=7.0Hz,2H), 4.97(s,2H), 7.10-7.35(m,5H)

Elemental analysis (C<sub>18</sub>H<sub>26</sub>N<sub>2</sub>O 0.6H<sub>2</sub>O)

Calcd.: C,72.74: H,9.22: N,9.43

Found : C,72.74: H,9.00: N,9.32

## I-65:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.25(s,18H), 1.31(d,J=7.0Hz,6H), 2.41(s,3H), 2.97(m,1H), 3.94(s,2H), 4.24(s,2H), 5.04(s,2H), 6.90(d,2H), 7.08-7.36(m,6H)

Elemental analysis (C <sub>30</sub> H <sub>42</sub> N <sub>2</sub> O 0.6H <sub>2</sub> O)			
Calcd.:	C,78.76:	H,9.52:	N,8.12
Found :	C,78.67:	H,9.31:	N,8.15

## I-66:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.04(t,J=7.0Hz,3H), 1.27(s,18H), 1.34(d,J=7.0Hz,6H), 2.50(s,3H), 3.21(q,J=7.0Hz,2H), 3.96(s,2H), 4.99(s,2H), 6.93(d,2H), 7.26(d,1H)

## I-67:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.08(t,J=7.0Hz,3H), 1.26 (d,J=7.2Hz,6H), 2.43(s,3H), 2.85(m,1H), 3.33(q,J=7.0Hz,2H), 3.94(s,2H), 4.94(s,2H), 6.98(d,J=1.6Hz,2H), 7.20 (d,J=1.6Hz,1H)

Elemental analysis (C <sub>17</sub> H <sub>22</sub> N <sub>2</sub> OCl <sub>2</sub> 0.2H <sub>2</sub> O)				
Calcd.:	C,59.20:	H,6.55:	N,8.12:	C,120.56
Found :	C,59.00:	H,6.52:	N,8.13:	C,120.58

## I-68:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.27(d,J=6.8Hz,6H), 2.40(s,3H), 2.86(m,1H), 3.88(s,2H), 4.37(s,2H), 4.96(s,2H), 6.87-7.34(m,8H)

## I-69:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.27(d,J=6.6Hz,6H), 2.39(s,3H), 2.88(m,1H), 3.91(s,2H), 4.38(s,2H), 4.95(s,2H), 6.50-7.22(m,8H)

## I-70:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.09(t,J=7.0Hz,3H), 1.26 (d,J=6.8Hz,6H), 2.43(s,3H), 2.86(m,1H), 3.34(q,J=7.0Hz,2H), 3.97(s,2H), 4.93(s,2H), 6.58-6.70(m,3H)

## I-88:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.26(d,J=7.2Hz,6H), 2.26(s,9H), 2.27(s,3H), 3.00(m,1H), 3.87(s,2H), 4.48(s,2H), 5.19(s,2H), 6.85(s,2H), 7.28-7.41(m,5H)

## I-89:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.19(t,J=7.0Hz,3H), 1.26 (d,J=7.2Hz,6H) 2.25(s,6H), 2.30(s,3H), 2.31(s,3H), 3.01(m,1H), 3.44(q,J=7.0Hz,2H), 3.85(s,2H), 5.13(s,2H), 6.83(s,2H)

## I-90:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.22(t,J=7.0Hz,3H), 1.24 (d,J=7.4Hz,6H), 2.41(s,3H), 3.06(m,1H), 3.48(q,J=7.0Hz,2H), 3.89(s,2H), 5.17(s,2H), 7.07-7.15(m,3H)

1-92

1-93:

10

15

## 20



35

40

45

50

55

55

I-94


<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.26(d,J=7.2Hz,6H), 2.40(s,3H), 3.08(m,1H), 3.75(s,6H), 3.91(s,2H), 4.50(s,2H), 5.22(s,2H), 6.25-6.45(m,3H), 7.25-7.42(m,5H)

Elemental analysis (C<sub>24</sub>H<sub>30</sub>N<sub>2</sub>O<sub>3</sub> 1.3H<sub>2</sub>O)

Calcd.(%);	C,68.97;	H,7.86;	N,6.70
Found (%);	C,68.89;	H,7.64;	N,6.52

## Example 72

5-(3,5-Dimethoxybenzyl)-1-ethoxymethyl-4-isopropyl-2-methylimidazole (I-72) and 4-(3,5-dimethoxybenzyl)-1-ethoxymethyl-5-isopropyl-2-methylimidazole (I-95)



## (1) Diethyl 3,5-dimethoxyacetamidomalonate 7

To a solution of 2.47 g of metallic sodium (107 mmol) in 247 ml of ethanol is added 23.28 g of diethyl acetamidomalonate (107 mmol). The resultant mixture is stirred at room temperature for 30 minutes, mixed with 20 g of 3,5-dimethoxybenzyl chloride (107 mmol), heated under reflux for 12 hours and concentrated in vacuo. The residue is extracted with ethyl acetate, washed with saturated brine, dried over sodium sulfate, and filtered. The filtrate is concentrated. The crude product is recrystallized from ethyl acetate/n-hexane to give 22.12 g of the objective diethyl 3,5-dimethoxybenzylacetamidomalonate 7. Yield, 56.18%.

7:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.30(t,J=7.4Hz,6H), 2.04(s,3H), 3.73(s,6H), 3.58(s,2H), 4.28(q,J=7.2Hz,4H), 6.15-6.35 (m,3H), 6.58(br,1H)

Elemental analysis (C <sub>18</sub> H <sub>25</sub> NO <sub>7</sub> )			
Calcd.(%);	C,58.85:	H,6.86:	N,3.81
Found (%);	C,58.67:	H,6.73:	N,3.82

(2) N-acetyl-3,5-dimethoxyphenylalanine 8

A mixture of 8 g of diethyl 3,5-dimethoxybenzylacetamidomalonate (21.8 mmol) and 10% aqueous sodium hydroxide is heated under reflux for 4 hours. After cooling, the reaction mixture is mixed with 17.74 ml of 36% hydrochloric acid and heated under reflux for 30 minutes. The mixture is allowed to cool, and the precipitated crystals are filtered and dried to give 5.30 g of the objective N-acetyl-3,5-dimethoxyphenylalanine 8. Yield, 91.06%.

8:<sup>1</sup>H-NMR(CD<sub>3</sub>OD-TMS)ppm: 1.91(s,3H), 3.00(m,2H), 3.74(s,6H), 4.65(m,1H), 6.30-6.40(m,3H)

Elemental analysis (C <sub>13</sub> H <sub>17</sub> NO <sub>5</sub> 0.3H <sub>2</sub> O)			
Calcd.(%);	C,57.26:	H,6.51:	N,5.14
Found (%);	C,57.09:	H,6.24:	N,5.13

(3) N-acetyl-3,5-dimethoxyphenylalanine methyl ester 9

To a mixture of 50 ml of 25% aqueous sodium hydroxide and 100 ml of ethyl ether is added 5 g of N-nitrosomethylurea at 10°C, and the mixture is stirred vigorously for 10 minutes. The ethyl ether layer is separated, dried over 15 g of potassium hydroxide and added to a solution of 5.3 g of N-acetyl-3,5-dimethoxyphenylalanine (19.8 mmol) in 100 ml of tetrahydrofuran, and the mixture is stirred at 10°C for 1 hour and concentrated in vacuo. The crude product is washed with ethyl ether to give 5.16 g of the objective N-acetyl-3,5-dimethoxyphenylalanine methyl ester 9. Yield, 92.51%.

9:<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 2.00(s,3H), 3.06(d,J=5.7Hz,2H), 3.75(s,9H), 4.87(m,1H), 5.92(br,1H), 6.23-6.36(m,3H)


Elemental analysis (C <sub>14</sub> H <sub>19</sub> NO <sub>6</sub> 0.2H <sub>2</sub> O)			
Calcd.(%);	C,59.02:	H,6.86:	N,4.92
Found (%);	C,58.91:	H,6.69:	N,5.07

The subsequent reactions were carried out in the same manner as in Example 42 to give the objective compounds (I-72) and (I-95).

I-72:<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)ppm: 1.11(t,J=7.0Hz,3H), 1.28(d,J=6.6Hz, 6H), 2.42(s,3H), 2.91(m,1H), 3.34(q,J=7.0Hz,2H), 3.74(s,6H) 3.92(s,2H) 4.93(s,2H) 6.22-6.30 (m,3H)Examples 73 - 85


In examples 73 to 85, compounds (I-73) - (I-85) as shown in the following Tables 9 and 10 were prepared substantially in accordance with the procedures used for the production of compound (I-86) in Example 31.

Table 9



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
73	S	-CH <sub>3</sub>		-CH <sub>2</sub>	-H	oil	1-73
74	S	-CH <sub>3</sub>		-CH <sub>2</sub>	-Br	oil	1-74
75	S	-CH <sub>3</sub>		-CH <sub>2</sub>	-COOEt	oil	1-75
76	S			-CH <sub>2</sub>	-H	mp 140-141°C (HCl salt)	1-76
77	S	-CH <sub>3</sub>		-C <sub>2</sub> H <sub>5</sub>	-H	oil	1-77
78	S	-CH <sub>3</sub>	-n-C <sub>4</sub> H <sub>9</sub>	-CH <sub>2</sub>	-H	oil	1-78
79	S	-C <sub>2</sub> H <sub>5</sub>		-CH <sub>2</sub>	-H	mp 90-91°C (HNO <sub>3</sub> salt)	1-79

Table 10



Exmple No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	mp(°C)	Compound No.
80	S	-H		-CH <sub>2</sub> -	-H	mp 93°C	1-80
81	S	-C <sub>2</sub> H <sub>5</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1-81
82	S	-C <sub>2</sub> H <sub>5</sub>		-CH <sub>2</sub> -	-CH <sub>3</sub>	oil	1-82
83	S	-C <sub>2</sub> H <sub>5</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	oil	1-83
84	S	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub> -	-CH <sub>3</sub>	oil	1-84
85	S	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub> -	-CH <sub>3</sub>	oil	1-85

Physical properties of the compounds shown in Tables 9 and 10 are provided below.

1-73:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.35(s,3H), 4.49(s,2H), 5.31(s,2H), 7.10-7.39(m,10H), 7.67(s,1H)

1-74:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 2.37(s,3H), 4.56(s,2H), 5.37(s,2H), 7.10-7.36(m,10H)

1-75:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.42(t,J=7.0Hz,2H), 2.39(s,3H), 4.42(q,J=7.0Hz,2H), 4.59(s,2H), 5.96(s,2H), 7.13-7.31 (m,10H)

1-76: (free base)

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 4.43(s,2H), 5.40(s,2H), 7.03-7.39 (m,13H), 7.95(s,1H), 8.06.11(m,2H)



I-77:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.20(t,J=7.0Hz,3H), 2.34(s,3H), 3.47(q,J=7.0Hz,2H), 5.26(s,2H), 7.09-7.26(m,5H), 7.67(s,1H)

I-78:

5 <sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 0.89(t,J=7.5Hz,3H), 1.47(m,4H), 2.32(s,3H), 2.79(t,J=7.5Hz,2H), 4.44(s,2H), 5.27(s,2H), 7.26-7.38(m,5H), 7.55(s,1H)

I-79: (free base)

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.13(t,J=7.5Hz,3H), 2.79 (q,J=7.5Hz,2H), 4.49(s,2H), 5.32(s,2H), 7.10-7.35(m,10H), 7.63(s,1H)

10 I-80:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 4.49(s,2H), 5.32(s,2H), 7.17-7.39 (m,11H), 7.72(s,1H)

I-81:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.11(t,J=7.0Hz,3H), 1.23 (t,J=7.0Hz,3H), 2.48(s,3H), 2.75(q,J=7.0Hz,2H), 3.49(q,J=7.0Hz,2H), 5.20(s,2H), 7.00-7.30(m,5H)

15 I-82:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.10(t,J=7.4Hz,3H), 2.20(s,6H), 2.42(s,3H), 2.73(q,J=7.4Hz,2H), 4.50(s,2H), 5.26(s,2H), 6.70-6.77(m,3H), 7.32-7.36(m,5H)

I-83:

20 <sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.12(t,J=7.6Hz,3H), 1.22 (t,J=7.0Hz,3H), 2.21(s,6H), 2.48(s,3H), 2.75(q,J=7.6Hz,2H), 3.50(q,J=7.0Hz,2H), 5.21(s,2H), 6.70.77 (m,3H)

I-84:

<sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.33(d,J=7.0Hz,6H), 2.20(s,6H), 2.40(s,3H), 3.19(sept,1H), 4.52(s,2H), 5.28(s,2H), 6.60(s,2H), 7.15-7.20(m,6H)

I-85:

25 <sup>1</sup>H-NMR(CDCl<sub>3</sub>-TMS)δppm: 1.33(d,J=7.0Hz,6H), 2.41(s,3H), 3.18(sept,1H), 4.54(s,2H), 5.28(s,2H), 7.00-7.37(m,10H)

Compounds (I) obtained in the foregoing Examples are evaluated as to the anti-retrovirus activity and the cytotoxicity.

30 Examples 86 - 137

In the same manner as described above, Compounds I-96 to I-147 are prepared.

35


40

45

50


55

Table 11




Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Compound No.
86	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-96
87	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-97
88	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-98
89	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-99
90	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-100
91	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-101
92	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-102
93	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-103
94	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-104
95	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-105
96	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-106
97	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-107
98	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-108

Table 12




Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Compound No.
99	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-109
100	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-110
101	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-111
102	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-112
103	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-113
104	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-114
105	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-115
106	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-116
107	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-117
108	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-118
109	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-119
110	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-120
111	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-121

Table 13



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Compound No.
112	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-122
113	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-123
114	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-124
115	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-125
116	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-126
117	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-127
118	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-128
119	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-129
120	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-130
121	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-131
122	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-132
123	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-133
124	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-CH <sub>2</sub>	-CH <sub>3</sub>	I-134

Table 14



Example No.	X	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	Compound No.
125	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-135
126	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-136
127	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-137
128	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-138
129	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-139
130	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-140
131	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-141
132	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-142
133	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-143
134	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-144
135	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-145
136	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-146
137	CH <sub>2</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>		-C <sub>2</sub> H <sub>5</sub>	-CH <sub>3</sub>	I-147

## Experiment 1

Infection Inhibitory Test

5 Anti-HIV activity and cytotoxicity are tested in the following manners.

1. Antivirus Activity

10 (1) A human T cell MOLT-4 clone 8, a strain continuously infected with HIV (HTLV-III<sub>B</sub> strain), was incubated in RPMI-1640 medium containing 10% bovine fetus serum and the supernatant was isolated by filtration. After the determination of virus titer, the supernatant was stored at -80°C. A compound to be tested was diluted with the same medium to a given concentration and poured into 96-well microtiter plates by 100 µl/well. To the well is added 50 µl (2.5 x 10<sup>4</sup> cells) MT-4 cell suspensions, followed by the addition of 50 µl [600 pfu (plaque forming unit)] HIV-containing supernatant previously diluted with the same medium.

15 (2) After 5-day-incubation at 37°C in a carbon dioxide incubator, 30 µl MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide), 5 mg/ml in PBS) was added to each well, and the incubation was continued for another 1 hour, during which surviving cells reduce MTT to precipitate formazan.

20 The resultant formazan was eluted by replacing 150 µl supernatant of a culture in each well by 150 µl solution comprising 10% triton X - 100 and isopropanol containing 0.4 v/v % HCl and shaking by means of a plate mixer. The resultant formazan was assayed at OD 540 nm, and the result was evaluated by comparison with that obtained in a control test.

IC<sub>50</sub> is defined as a concentration of the compound (I) that gives a 50% inhibition against viral cytotoxicity. The results are summarized in Table 15.

Table 15



Compound No.	IC <sub>50</sub> (µg/ml)	Compound No.	IC <sub>50</sub> (µg/ml)
I-1	0.03-0.08	I-50	0.016
I-4	0.13-0.25	I-58	0.5
I-5	0.13-0.25	I-67	0.005
I-12	0.06-0.13	I-69	0.003
I-14	0.016-0.032	I-90	0.5-1.0
I-15	0.063-0.12	I-91	0.5-1.0
I-16	0.016-0.031	I-92	0.25
I-17	0.016	I-93	0.06
I-18	0.016	AZT	0.005-0.01
I-29	0.13-0.25	TIBO(R-82913)	0.13-0.25
I-42	0.016		

2. Cytotoxicity

50 Cytotoxicity of a compound (I) was evaluated in a similar manner as described in 1 above except that 50 µl of the same medium is added to each well in place of the HIV-containing supernatant (viral solution).

CC<sub>50</sub> is defined as a concentration of the compound (I) that gives 50% cytotoxicity.

Results are summarized in Table 16 below.

Table 16


Compound No.	CC <sub>50</sub> (μg/ml)	Compound No.	CC <sub>50</sub> (μg/ml)
I - 4	12.5 - 25	I - 50	6.25 - 12.5
I - 5	12.5 - 25	I - 58	> 100
I - 9	12.5 - 25	I - 67	6.25 - 12.5
I - 12	12.5 - 25	I - 69	6.25 - 12.5
I - 13	12.5 - 25	I - 90	1.0 - 10
I - 14	10 - 20	I - 91	10 - 100
I - 15	10 - 20	I - 92	6.25 - 12.5
I - 18	25 - 50	I - 93	12.5 - 25
I - 29	25 - 50	A Z T	25
I - 42	3.1 - 6.25	TIBO(R-82913)	12.5 - 25

As is clear from the results of Tables 11 and 12, the compounds of the present invention or salts thereof possess anti-HIV activity, whereby the compounds of the invention can be used for the treatment of HIV virus infections such as AIDS or the like but also the development of derivatives improved pharmacological properties.

In the compounds of formula I the various alkyl groups may be substituted provided that the resulting compound has anti-viral activity.


#### Claims

1. A compound of the formula I:



wherein R<sup>1</sup> is hydrogen, alkyl, halogen, or optionally substituted aryl, R<sup>2</sup> is alkyl, optionally substituted aryl optionally substituted aralkyl, or an optionally substituted hetero ring group; R<sup>3</sup> is hydrogen, alkyl, optionally substituted aryl, optionally substituted aralkyl, or optionally substituted hydroxyalkyl; R<sup>4</sup> is hydrogen, alkyl, halogen, acyl, optionally substituted hydroxyalkyl, optionally esterified or amidated carboxyl group, hydroxy group, aryl or arylthio; X is S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or Se; n is an integer of 1 to 3, or a pharmaceutically acceptable salt thereof.

2. A compound according to Claim 1, wherein R<sup>1</sup> is hydrogen, lower alkyl, halogen, or phenyl; R<sup>2</sup> is lower alkyl, phenyl optionally substituted by lower alkyl or halogen or pyridyl; R<sup>3</sup> is lower alkyl, benzyl, or hydroxyalkyl; R<sup>4</sup> is hydrogen, lower alkyl, halogen, formyl, optionally substituted hydroxymethyl, optionally esterified or amidated carboxyl group, hydroxy group or phenylthio group; and X is S or CH<sub>2</sub>; and n is a integer of 1.
3. A pharmaceutical composition comprising an effective amount of a compound of Claim 1 or Claim 2 in association with a pharmaceutically acceptable, substantially nontoxic carrier or excipient.
4. An antiviral agent which comprises an effective amount of a compound of Claim 1 or Claim 2 together with pharmaceutically acceptable carriers therefor.
5. The use of a compound of the formula I:



wherein R<sup>1</sup> is hydrogen, alkyl, halogen, or optionally substituted aryl, R<sup>2</sup> is alkyl, optionally substituted aryl optionally substituted aralkyl, or an optionally substituted hetero ring group; R<sup>3</sup> is hydrogen, alkyl, optionally substituted aryl, optionally substituted aralkyl, or optionally substituted hydroxyalkyl; R<sup>4</sup> is hydrogen, alkyl, halogen, acyl, optionally substituted hydroxyalkyl, optionally esterified or amidated carboxyl group, hydroxy group, aryl or arylthio; X is S, SO, SO<sub>2</sub>, CH<sub>2</sub>, or Se; n is an integer of 1 to 3, or a pharmaceutically acceptable salt thereof, in the preparation of a pharmaceutical formulation for the treatment of a retrovirus infection, such as human immunodeficiency syndromes.





European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 93 30 0278

Page 1

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	JP-A-63 150 266 (MITSUI PETROCHEMICAL INDUSTRIES, LTD.; SUNTORY, LTD.) 22 June 1988 * 5-[(2,6-dimethylphenyl)methyl]-1H-imidazole-1-methanol *	1,2	C07D233/84 C07D233/90 C07D233/60 C07D401/06 C07D403/06 C07D413/06 C07D417/06
P,X	EP-A-0 477 778 (EISAI CO., LTD.) 1 April 1992 * entire document *	1,2	A61K31/35 A61K31/415 A61K31/395
X	J. CHEM. SOC., PERKIN TRANS. 1 no. 7, 1987, pages 1445 - 1451 B. IDDON, N. KHAN 'Azoles. Part 5. Metal-halogen exchange reactions of polybromoimidazoles' * 2,4-dibromo-1-(methoxymethyl)-5-(methylthio)-1H-imidazole *	1,2	A61K31/42 A61K31/425 A61K31/435 A61K31/44 A61K31/495 A61K31/50 A61K31/505 C07D233/54
X	J. CHEM. SOC., PERKIN TRANS. 1 no. 4, 1983, pages 735 - 738 B. IDDON, B.L. LIM 'Metal-halogen exchange reactions of mono- and polyhaloimidazoles' * 1-(ethoxymethyl)-4-iodo-2,5-bis(phenylthio)-1H-imidazole *	1,2	TECHNICAL FIELDS SEARCHED (Int. Cl.5)  C07D A61K
X	J. CHEM. SOC., PERKIN TRANS. 1 no. 2, 1983, pages 279 - 283 B. IDDON, B.L. LIM 'Metalation and metal-halogen exchange reactions of imidazoles' * 1-(ethoxymethyl)-2,5-bis(phenylthio)-1H-imidazole; 1-(ethoxymethyl)-5-(methylthio)-2-(phenylthio)-1H-imidazole-4-carboxaldehyde *	1,2	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 16 APRIL 1993	Examiner HERZ C.P.
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1503 (01.87) (PMA)



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 93 30 0278

Page 2

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
X	J. CHEM. SOC., CHEM. COMMUN. no. 21, 1981, pages 1095 - 1097 B. IDDON, B.L. LIM 'Metalation and metal-halogen exchange reactions of imidazoles' * compounds of formula I *	1,2	
A	CHEM. PHARM. BULL. vol. 39, no. 4, 1991, pages 871 - 875 C. KANEKO ET AL. 'A series of novel acyclic nucleosides. IV. Synthesis of N(1)-sulfur analogs of acyclovir, directed toward improved antiviral activities'	1-5	
A	NUCLEOSIDES NUCLEOTIDES vol. 9, no. 6, 1990, pages 815 - 826 D.P.C. MCGEE ET AL. 'Synthesis and antiviral activity of the 3-deaza analog of 9-[(1,3-dihydroxy-2-propoxy)methyl]guanine'	1-5	
A	J. MED. CHEM. vol. 28, no. 6, 1985, pages 834 - 838 R. ALONSO ET AL. 'Synthesis and antiviral evaluation of nucleosides of 5-methylimidazole-4-carboxamide'	1-5	
The present search report has been drawn up for all claims			
Place of search MUNICH		Date of completion of the search 16 APRIL 1993	Examiner HERZ C.P.
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons A : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			

RPO FORM 150 (12/92) (P0001)